



Market Areas



● Registered Wells > 500 GPM

| | | | | | |
|------|------|------|------|------|------|
| 2439 | 2437 | 2435 | 2433 | 2431 | 2429 |
| 2607 | 2609 | 2611 | 2613 | 2615 | 2617 |
| 2733 | 2731 | 2729 | 2727 | 2725 | 2723 |
| 2903 | 2905 | 2907 | 2909 | 2911 | 2913 |
| 3029 | 3027 | 3025 | 3023 | 3021 | 3019 |
| 3199 | 3201 | 3203 | 3205 | 3207 | |



Legend

- Towns
- Sections
- Rivers and Streams
- Topography

Soil Classes

- Lakes and Ponds
- Excessively drained sandy soils formed in alluvium in valleys and eolian sand on uplands in sandhills
- Excessively drained sandy soils formed in eolian sands on uplands in sandhills
- Moderately well drained silty soils on uplands and in depressions formed in loess
- Well drained silty soils formed in loess on uplands
- Well drained silty soils formed in loess and alluvium on stream terraces
- Well to somewhat excessively drained loamy soils formed in weathered sandstone and eolian material on uplands
- Somewhat poorly drained soils formed in alluvium on bottom lands
- Moderately well drained silty soils with clayey subsoils on uplands

Sherman County