



Exhibit 85 A - page 1



● Registered Wells > 500 GPM

|      |      |      |      |      |      |
|------|------|------|------|------|------|
| 3993 | 3991 | 3989 | 3987 | 3985 | 3983 |
| 4143 | 4145 | 4147 | 4149 | 4151 | 4153 |
| 4233 | 4231 | 4229 | 4227 | 4225 | 4223 |
| 4383 | 4385 | 4387 | 4389 | 4391 | 4393 |
| 4479 | 4477 | 4475 | 4473 | 4471 | 4469 |
|      |      |      |      |      |      |



### Legend

- Towns
- Sections
- Rivers and Streams
- Topography

### Soil Classes

- Lakes and Ponds
- Excessively drained sandy soils formed in alluvium in valleys and eolian sand on uplands in sandhills
- Excessively drained sandy soils formed in eolian sands on uplands in sandhills
- Moderately well drained silty soils on uplands and in depressions formed in loess
- Well drained silty soils formed in loess on uplands
- Well drained silty soils formed in loess and alluvium on stream terraces
- Well to somewhat excessively drained loamy soils formed in weathered sandstone and eolian material on uplands
- Somewhat poorly drained soils formed in alluvium on bottom lands
- Moderately well drained silty soils with clayey subsoils on uplands